Hua Loo Keng's Problem Involving Primes of a Special Type

Sergey A. Gritsenko and Natalya N. Motkina

arXiv:0812.4665v1 [math.NT] 26 Dec 2008

Abstract

Let η be a quadratic irrationality. The variant of Hua Loo Keng's problem involving primes such that $a < \{\eta p^2\} < b$, where a and b are arbitrary real numbers of the interval $(0,1)$, solved in this paper.

1. Introduction Suppose that $k \geq 2$ and $n \geq 1$ are positive integers. Consider the equation

$$p_1^n + p_2^n + \dots + p_k^n = N \tag{1}$$

in primes $p_1,\ p_2, \dots, p_k$.

Several classical problems in number theory can be reduced to the question on the number of solutions of the equation (1). For example, if $n = 1$ and $k = 2$ or 3 then (1) is the equation of Goldbach; if $n \geq 3$ then (1) is the equation of Waring–Goldbach.

Let $\mathcal{P}$ be a subset of the set of primes, $\mu = \lim\limits_{n\to\infty} \pi^{-1}(N) \sum\limits_{\substack{p\leq N\\ p\in\mathcal{P}}} 1$ be the "density" of $\mathcal{P}$, $0 < \mu < 1$.

Let $J_{k,n}(N)$ be the number of solutions of (1) in primes of the set $\mathcal{P}$, and $I_{k,n}(N)$ be the number of solutions of (1) in arbitrary primes.

Sometimes the equality

$$J_{k,n}(N) \sim \mu^k I_{k,n}(N) \tag{2}$$

holds.

For example, if $\mathcal{P} = \{p \mid a < \{\sqrt{p}\} < b\}$, $0 < a < b < 1$ then provided $n = 1$, $k = 3$, and also provided $n \geq 3$, $k \gg n^2 \log n$ the equality (2) holds (s. [1]).

In present paper we solve the additive problem for which the property (2) does not hold.

Let η be a quadratic irrationality, $0 < a < b < 1$, $\mathcal{P} = \{p \mid a < \{\eta p^2\} < b\}$, $n = 1$. Then $\mu = b - a$ (s. [2]).

The main result of the present paper is the following theorem which shows that the equality

$$J_{5,2}(N) \sim (b-a)^5 I_{5,2}(N) \tag{3}$$

does not hold.

Note that (s. [3])

$$I_{5,2}(N) \asymp \frac{N^{3/2}}{\log^5 N}$$

provided $N \equiv 5(\mod 24)$.

Theorem 1. *Suppose*

$$\sigma(N,a,b) = \sum_{|m|<\infty} e^{2\pi i m(\eta N - 2.5(a+b))} \frac{\sin^5 \pi m(b-a)}{\pi^5 m^5}.$$

Then the following formula holds

$$J_{5,2}(N) = I_{5,2}(N)\sigma(N,a,b) + O(N^{3/2-0,00002}).$$

Outline of proof. Define the function

$$\psi_0(x) = \begin{cases} 1, & \textit{if} \quad a < x < b, \\ 0, & \textit{if} \quad 0 \le x \le a \quad \textit{or} \quad b \le x \le 1 \end{cases}$$

and extend it by periodicity to the entire numerical axis. So that

$$J_{5,2}(N) = \int_0^1 \left(\sum_{p \le \sqrt{N}} \psi_0(\eta p^2) e^{2\pi i x p^2} \right)^5 e^{-2\pi i x N} dx.$$

We record Vinogradov's lemma (see [1], Section III, Chapter 2, Lemma 2). In the notation of the lemma, let $r = [\ln N]$, $\Delta = N^{-0,01}$. Denote by ψ_1 a function ψ from the lemma with $\alpha = a + \Delta/2$, $\beta = b - \Delta/2$ and by ψ_2 with $\alpha = a - \Delta/2$, $\beta = b + \Delta/2$. Then we have the inequality

$$J_1(N) \le J_{5,2}(N) \le J_2(N), \tag{4}$$

where

$$J_k(N)=\int_0^1\left(\sum_{p\le\sqrt{N}}\psi_k(\eta p^2)e^{2\pi ixp^2}\right)^5 e^{-2\pi ixN}dx,\quad k=1,2.$$

Next we derive an asymptotic formulas for $J_1(N)$ and $J_2(N)$. The main terms of these formulas coincide. Use the Fourier series for $\psi_k(x)$

$$\psi_k(\eta p^2)=\sum_{|m|\le r\Delta^{-1}}c_k(m)e^{2\pi im\eta p^2}+O(N^{-\ln\pi}),$$

thus

$$J_k(N)=\sum_{|m_1|\le r\Delta^{-1}}c_k(m_1)\sum_{|m_2|\le r\Delta^{-1}}c_k(m_2)\cdot$$

$$\cdot\sum_{|m_3|\le r\Delta^{-1}}c_k(m_3)\sum_{|m_4|\le r\Delta^{-1}}c_k(m_4)\sum_{|m_5|\le r\Delta^{-1}}c_k(m_5)\cdot$$

$$\cdot\int_0^1 S(x+m_1\eta)S(x+m_2\eta)S(x+m_3\eta)S(x+m_4\eta)S(x+m_5\eta)e^{-2\pi ixN}dx$$

$$+O(N^{3/2-\ln\pi}\ln N),$$

where

$$S(x)=\sum_{p\le\sqrt{N}}e^{2\pi ixp^2}.$$

Note that

$$\sum_{|m|\le r\Delta^{-1}}c_k^5(m)\int_0^1 S^5(x+m\eta)e^{-2\pi ixN}dx=$$

$$=I_{5,2}(N)(\sigma(N,a,b)+O(\Delta)).\tag{5}$$

Consider the sets such that $(m_1,m_2,m_3,m_4,m_5)\ne(m,m,m,m,m)$ and the integrals

$$I(N,m_1,m_2,m_3,m_4,m_5)=$$

$$=\int_0^1|S(x+m_1\eta)||S(x+m_2\eta)||S(x+m_3\eta)||S(x+m_4\eta)||S(x+m_5\eta)|dx.$$

Without loss of generality it can be assumed that $m_1 < m_2$. By t denote $x + m_1\eta$. Since the integrand has a period 1, we can take t to lie in the range $E = [-1/\tau; 1 - 1/\tau)$, where $\tau = N^{1-0,001}$. Suppose $d, q \in \mathbb{Z}$ are such that

$$t = \frac{d}{q} + \frac{\theta_1}{q\tau}, \quad (d,q) = 1, \quad 1 \le q \le \tau, \quad |\theta_1| < 1.$$

Divide the interval E into two intervals E_1 and E_2 for which $q \le N^{0,001}$ and $N^{0,001} < q \le \tau$ respectively. On writing $m_2' = m_2 - m_1$, ..., $m_5' = m_5 - m_1$, we can express the above integral as

$$\int_{-1/\tau}^{1-1/\tau} |S(t)||S(t+m_2'\eta)||S(t+m_3'\eta)||S(t+m_4'\eta)||S(t+m_5'\eta)|dt =$$

$$= \int_{E_1} F(t)dt + \int_{E_2} F(t)dt,$$

where

$$F(t) = |S(t)||S(t+m_2'\eta)||S(t+m_3'\eta)||S(t+m_4'\eta)||S(t+m_5'\eta)|.$$

It is well known that if $t \in E_2$ then

$$|S(t)| \ll N^{0,5-0,00002}.$$

Consider $|S(t+m_2'\eta)|$ when $t \in E_1$. Since η is the quadratic irrationality then we have

$$t + m_2'\eta = \frac{X}{Y} + \frac{\theta_2}{Y^2}, \quad (X,Y) = 1, \quad |\theta_2| < N^{0,012}\ln N \qquad (6)$$

and

$$\sqrt{\tau}N^{-0,01015}\ln N < Y < \sqrt{\tau}N^{0.001}. \qquad (7)$$

Then from (6) and (7) it follows in the standard way the estimate

$$|S(t+m_2'\eta)| \ll N^{0,5-0,00002}.$$

Using the inequality between arithmetical and geometric means, we finally obtain

$$I(N, m_1, m_2, m_3, m_4, m_5) \ll N^{3/2-0,00002}. \qquad (8)$$

Now the theorem follows directly from formulas (4), (5), (8).

Remark. If $\mathcal{P} = \{p \mid a < \{\eta p\} < b\}$ then the equality (3) holds.

References

[1] S.A. Gritsenko, *Three additive problems*, Izvestiya Mathematics, 1993, 41 (3), 447–464

[2] I.M.Vinogradov, *Special variant of the method of trigonometric Sums*, Nauka, Moscow 1976; In: *Selected works*, Springer, Berlin, 1985.

[3] I.M.Vinogradov, *Method of trigonometric Sums in number theory* [in Russian], Nauka, Moscow 1971.